

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

5 Feet, or 11 Paris Miles (at 2000 Toises the Mile) 232 Toises, 5 Foot. But as in order to determine the whole Height of the Atmosphere, the Logarithm of 1" ought to be deducted from the Logarithm of 336" or 28" o", and as that Logarithm is 00000, it follows from thence, that beyond the Place, where the Mercury would descend to 1", the Air is expanded into an Indefinite Space.

For the Satisfaction of the Curious, I have added the Tables themselves, to wit, those which Cassini the Younger calculated according to the Rules of Mariotte, those which he deduced from the Observations made by the Gentlemen of the Royal Academy of Sciences, who drew the Meridian Line, and those which my Uncle calculated from the Observation made

at Pfeffers in 1709.

In another Paper on this Subject I intend to compare the Height of Mountains, as determined by divers antient and modern Writers, with the true Height of them, as it appears to be by the Barometrical Observations, particularly those made by my Father on the high Mountains of Swifferland.

SIR,

A FTER I had mentioned my being pretty well affured, that the Plants of Misleto were some of them Male, and some Female; and had promised to D d d d 2 com-

II. Observations of a Difference of Sex in Misseto, in a Letter from the Reverend Mr. Edmund Barrel to Sir Hans Sloane, Bart. &c.

communicate to you, such farther Observations as I should make of that Matter; I was discouraged from giving you any Trouble about it, by an Information I received, that the eminently learned Dr. Herman Boerhave had already told the World of a Difference of Sex in Milleto. But having, fince that, feen the Historia Plantarum, which is published as of his dictating; I find, that he mentions it in such a Manner as makes me suspect, that he only took his Notion from Tournefort, and was not fully apprized of the true Nature of these Plants. Ovarium alio à Flore loco natum feems to suppose both Flower and Ovary to be on the fame Plant, though in distinct Places of that Plant. I have therefore now written down my Observations for your Perusal in the same Order that I made them.

I have (from my own fowing of the Berries) four thriving Plants of Misleto growing on one Tree in my Garden. These, being often in my View, gave me the first Apprehension, of there being any Difference of Kind, or Sex, in this Shrub. They were not of Age to bear Flower or Fruit till 1726; when one of them bore a Berry or two; and expecting that they should all do so the following Year, I frequently examined them, and found that two Plants had Berries, and two had none. I then went and examined the Misleto on other Trees, which have Plants of above 20 Years Growth. And I find the Method of Nature to be thus.

Dr. Grew observes, that many Plants make a visible Preparation in the former Year for the Flower and Fruit of the next Season. This is done by Misleto. At the latter End of May, the Male Plants put out little

little Knobs, at the Joints and Tops of their Boughs; which at first are not very unlike the young green Berries; but they foon appear evidently distinct from them, and being by the latter End of July, grown as large as the Berries, are then not at all like them; foreading wider upwards, and having 3, or 4, or 5 Buds, at the Top of each Knob. About June, the Female Plant also makes a like Preparation; putting out at the Joints and Tops of the Boughs, Knobs, which are more sharp, and shorter than those of the Male; with 1, or 2, but most commonly with 3 Buds, or finall Points at the Top of each Knob. I call them Buds, because in their Season they open into Flowers, both in the Male and Female Plants; all the rest of the Knob serving only for Footstalks to the Flowers, in the one Sort, and to both Flower and Fruit in the other. By the latter End of August the Berries are grown much larger than the Knobs on the Male Plants. And from thence, till late in January, there is little worth Remark in either Plant; only the Berry grows fomewhat bigger, and becomes ripe; and the Knobs on the Male grow more and more yellow; fo that one may, at that Time, difcern a Male from a Female Plant, at a confiderable Distance. By the 20th of February Misses is in Bloom, both Male and Female. The Knobs of the Male are open at the Top with 3, or 4, or 5 Blossoms; which are very well described (though in short) in Boerhave's Historia Plantarum.

The Female Plant flowereth also now, with a Bloffom (which *Boerhave* calls the Ovarium) exactly like the Male Flower, save only, that the whole Female Flower is not bigger than one Leaf of the Male Flower. Flower. They both continue in full Bloom till the Middle of *March*, when the Male Blossoms begin to wither and drop off. And by the 20th of *March* the young Berries begin to shew themselves, swelling forth, one under each Female Blossom; which often adheres to the Top of the Berry; and being carried up with it, presently withers, and soon falls off again; tho' some continued on till the 12th of *May*, when the Berries were of the Size of a great Pin's Head.

This compleated the Year's Observation. And I think it is much to be wondered at, that this Plant. which hath been the Admiration of all Ages, should (scarce ever) find one Observer so curious as to follow the Changes of it, through one whole Year's Revolution. For if this had been done with any Accuracy, it must have been very evident, that one Sort of Milleto was very different from the other: One Sort bearing very small Flowers, with Berries succeeding them: the other bearing much larger Flowers, not fucceeded by any Berries; the very Footstalk of the Male falling off with the Flower; whereas the Footstalk of the Female, becomes a Footstalk to the Berry. It is possible, that this Difference of the Sex, in the Plants of Misseto, may be of Consequence in Medicine: To those therefore, who would make any Experiments of the different Virtues of these Plants, I offer this general Observation: That there is no Time of the Year wherein the Difference of these two Sorts, or Sexes of Misleto, is not very easy to be seen and known, by the Marks I have mentioned: And the meanest Herb-woman will foon have Skill enough to bring the Sort they are ordered to procure; there being as

great

great a Plenty of the Male Plants as there is of the Female. I follow your Commands, rather than my own Inclinations, in giving you this Trouble, and am, with the greatest Respect,

SIR,

Your most obliged, and
Sutton, near Dartford,
August 20, 1728. most obedient humble Servant,

Edmund Barrell.

III. An uncommon Sinking of the Ground in Kent. Communicated in a Letter to Mr. Peter Collinson.

SIR,

I T is within this three or four Days, that I have first been able to get a more particular Account of the finking of the Lands at Lymne in Kent, and even now, perhaps, it will be but an imperfect one.

It is now about two Years fince it happened, and was the Consequence of a very wet Season, when the Waters, that had fallen on the Up lands, and were not carried off by Drains, soaked into the Ground in such Quantities as to form a quick Sand at some considerable Depth in the Earth (at least this is what we look on to have caused the Phænomenon) which not being able to bear the Weight upon it, broke out at the Side of the Hill.